


1. (a) Draw a labelled diagram showing the atomic structure of $^{24}_{12}\text{Mg}$. (2 marks)



Nuclear Composition
 12p ✓
 12n ✓
 Electrons ✓



OR
 Energy levels ✓
 Nuclear with occupied protons and neutrons ✓

(b) The atomic number of phosphorus is 15. Draw a dot (•) and cross (x) diagram for the compound formed when phosphorus reacts with chlorine, atomic number 17. (1 mark)

1 or 0 marks
 - Use of only dots or crosses ✓
 - Crosses ✓




2. (a) State the condition under which a Bunsen burner produces a luminous flame. (1 mark)

When airhole/collar is closed ✓

fully closed.

Write an equation for the reaction that takes place in a luminous flame assuming the laboratory gas is butane. (1 mark)



(c) One of the regions in the non-luminous flame is the unburnt gas region. Describe how the presence of this region can be shown using a wooden splint. (1 mark)


Slip a wooden splint across the middle

Part of the flame

The central part remains unburnt/unchanged

while the outer part burns.

OR. unburnt region.



3. (a) The elements sodium, magnesium and aluminium belong to group I, II and III respectively. Select the element with the highest electrical conductivity and give a reason. (1 mark)

Aluminium $\checkmark \frac{1}{2}$

It has 3 delocalised $\checkmark \frac{1}{2}$ electrons unlike

Sodium and magnesium has one and two respectively.

- (b) Complete Table 1 to show the products of electrolysis for concentrated sodium chloride and molten sodium chloride.

Table 1

Compound	Anode	Cathode
Concentrated sodium chloride	Chlorine Cl_2	Hydrogen H_2
Molten sodium chloride	Chlorine Cl_2	Sodium Na

- Use of equation to show correct product Award!

(2 marks)

4. A small piece of sodium metal was placed in a beaker containing pure water.

- (a) State two observations made during the reaction.

(1 mark)

Two observations: The piece of metal starts floating

- melts into silvery ball.

- Production of effervescence / hissing sound

- The beaker becomes warm

- (b) State and explain another observation made when a drop of phenolphthalein is added to the mixture in the beaker. (1 mark)

Solution turns pink because sodium hydroxide / alkaline solution is formed.

Hydroxide / alkaline solution is formed.

- (c) Explain why it is not advisable to carry out this experiment using potassium metal. (1 mark)

Potassium reacts explosively with

water / more vigorously / more violent.

PDF Compressor Free Version


PDF Compressor Free Version

Stop marking
Candidate's name goes wrong
Use equations
Accepted partially

5. Describe how a pure sample of copper(II) nitrate crystals can be prepared using recycled copper wire.
Concentrated nitric acid or nitric acid
50% (3 marks)
acid

- Add excess unreacted copper
 - Filter the solution to saturation
 - Heat the solution to cool
 - Allow to form crystals
 - Dry/filter the crystals
6. The following apparatus and chemicals are used to investigate the percentage of air used when iron rusts: iron filings, 100 ml measuring cylinder, trough and water.

(a) Draw a setup of the experiment. (2 marks)



(b) Write an expression to show how the percentage of air used is calculated at the end of the experiment. (1 mark)

$$\frac{\text{Initial height of air column} - \text{Final height of air column}}{\text{Initial height of air column}} \times 100$$

OR

$$\frac{\text{Initial height of water} - \text{Final height of water}}{\text{Initial height of water}} \times 100$$

Volume instead of height

7. Figure 1 shows a graph of atomic radius of some group I and group II elements.




Figure 1

PDF Compressor Free Version

- (a) Explain why the atomic radius of sodium is higher than that of:

- (i) lithium.

$$Na = 2.8.1$$

$$Li = 2.1$$

(1 mark)

Sodium has 3 energy levels unlike Lithium

has two ✓ 1 or Li = 2.1 ✓ 1/2
Na = 2.8.1 ✓ 1/2

(ii) magnesium. $Mg = 2, 8, 2$ (1 mark)
 $Na = 2, 8, 1$ The effective

nuclear charge is higher in magnesium than.

Sodium has higher number of protons.

(b) Predict the atomic radius of calcium. (1 mark)


208 + 2 ✓ write out showing ✓ 1
✓ Extrapolate the value ✓ 1/2.

8. Compound D with formula, C_3H_4 , was reacted with excess hydrogen chloride gas. (1 mark)

(a) Give the name of compound D. (1 mark)

Propyne Prop-1-yne
Prop-1,2-diene

(b) Draw two possible structures of the products formed. (2 marks)



9. Study the setup in Figure 2 and answer the questions that follow.




Figure 2

PDF Compressor Free Version

- (a) State the precaution that should be taken in carrying out the experiment. Give a reason. (1 mark)

The experiment should be carried out in a fume chamber ^{1/2 out in open} since carbon (II) oxide is poisonous ^{1/2}

- (b) State the observations made in the boiling tube. (2 marks)

A white precipitate is formed which dissolves ^{1/2} to form a colourless solution.

10. Consider the following reaction:



The enthalpy change is -92.4 kJ per mole of nitrogen.

- (a) Give the enthalpy change per mole of ammonia. (1 mark)
- missing penultimate 1/2*
 $-\frac{92.4}{2} \Rightarrow -46.2 \text{ kJ mol}^{-1}$ ^{1/2} **11**
- (b) State and explain how each of the following affects the yield of ammonia: (1 mark)

- (i) Increase in temperature. (1 mark)

^{1/2} It lowers the yield of ammonia since

the forward reaction is exothermic ^{1/2}.

- (ii) Finely divided iron. (1 mark)

or backward reaction is endothermic.

No effect ^{1/2} A catalyst has no

effect on the position of the equilibrium ^{1/2}.

11. Study the flow chart in Figure 3 and answer the questions that follow.





Figure 3



(a) Identify solid E. $KMnO_4$ or MnO_4^- or PbO_2 . (1 mark)

(b) Name the type of reaction that takes place in chamber 1.
Potassium manganate(VII) or Manganese(IV) oxide
Lead(IV) oxide
Redox / oxidation (1 mark)

(c) Write an equation for the reaction that takes place in chamber 2. (1 mark)



12. Compounds H and J have the following structures.



(a) Give the names of:

(i) Compound H. (1 mark)

3-methylpentan-1-ol

(ii) Compound J.

(1 mark)

Butanoic acid.



(b) State the conditions necessary for H and J to react.

(1 mark)

Concentrated Sulphuric acid ✓
Warm / Heat ✓
Temperature between 30-60°C ✓


13. Rhombic sulphur is one of the allotropes of sulphur.

(a) Draw the structure of rhombic sulphur.


(1 mark)



OR
Octahedral
Structure



OR



• Sulphur.

(b)

Describe the observations made when rhombic sulphur is heated from room temperature until it boils.

(1 mark)

Yellow solid forms amber liquid ✓

(2 marks)


As the temp increases, the liquid becomes darker and more viscous ✓

Then it turns dark red brown and less viscous ✓

(1 mark)

14. The molar enthalpy of solution for potassium sulphate (K_2SO_4) is $+23.8 \text{ kJ}$.

- (a) On the axes provided, draw a labelled energy level diagram for the dissolution process of potassium sulphate in water. (2 marks)



(1 1/2 marks)


(b) Calculate the enthalpy change when 5.22 g of potassium sulphate is completely dissolved in water ($K = 39.0$; $S = 32.0$; $O = 16.0$). (2 marks)

R_{PM} of $K_2SO_4 = 174 \sqrt{1/2}$ (1 1/2 marks)

Moles of $K_2SO_4 = \frac{5.22}{174} = 0.03 \sqrt{1/2}$

$\Delta H = 0.03 \times 23.8 = 0.714 \text{ kJ} \sqrt{1/2}$ Wrong unit - p

For exact termic
rxn - penalise
fully.



PDF Compressor Free Version

15. (a) State Gay-Lussac's law.


(1 mark)

When gases react, they do so in volumes that bear simple ratios to one another and to the products if gaseous at constant temperature and pressure. ✓ 1

(b) 180 cm³ of nitrogen(II) oxide gas was reacted with 400 cm³ of oxygen gas.


(i) Write an equation for the reaction.

(1 mark)



(ii) Calculate the total volume of the gases at the end of the reaction.

(2 marks)



Using ratio:

$$\text{Volume of oxygen} = \frac{180 \times 1}{2} \quad \checkmark 1$$

$$= 90 \text{ cm}^3 \quad \checkmark 1$$

$$\text{Volume of oxygen unreacted} = 400 - 90 = 310 \quad \checkmark 1$$

$$\text{Volume of NO}_2 = 180 \text{ cm}^3 \quad \checkmark 1$$

$$\text{Total volume} = 310 + 180$$

$$= 490 \text{ cm}^3 \quad \checkmark 1$$

PDF Compressor Free Version

16.

Describe how the setup in Figure 4 can be used to distinguish between 50.0cm^3 of 0.2M hydrochloric acid and 50.0cm^3 of 0.2M ethanoic acid using pieces of 6m length of magnesium ribbon and a stop watch.

(3 marks)




Figure 4

- Any Contact between
Marking Points
- Mg and hydrochloric acid ✓
 - Mg and ethanoic acid ✓
 - Using a stop watch to show the difference ✓
 - Conclusion - HCl takes a shorter time ✓

- Put a 6cm Mg ribbon in conical flask and add 50cm^3 of HCl. Using a stop watch, record the volume of gases collected at a time interval every 15cm^3

- Repeat the experiment using 50cm^3 of ethanoic acid

- More Higher volume of gas will be collected when HCl is used than ethanoic acid at same interval of time

OR

The reaction rate will take a shorter time to completion when HCl is used than when ethanoic acid is used.

2023

PDF Compressor Free Version



17. Describe how dilute nitric(V) acid and blue litmus papers can be used to distinguish between solid samples of sodium carbonate and sodium sulphite. (3 marks)

To two different test tubes containing the samples
 Add dil. nitric(V) acid ✓₂
 - Place wet blue litmus test paper ✓₂ in test tubes
 - Both turn red ✓₂
 - There after one of them is bleached ✓₂
 - The sample that produces bleaching on the litmus is sodium sulphite ✓₁

18. (a) Describe how propanone can be used to extract a pure sample of sunflower oil. (2 marks)

Crush the sunflower seeds using motor and pestle ✓₂
 - Add propanone and stir ✓₂
 - Decant ✓₂
 - Leave the extract on sunlight for propanone to evaporate leaving oil behind.
 State why sodium hydroxide solution is not suitable for the extraction of sunflower oil. (1 mark)

It will react with oil to form soap ✓₁



19. 31.5 cm³ of concentrated nitric(V) acid was diluted to 500 cm³. 10.0 cm³ of the dilute acid required 25.0 cm³ of 0.4 M sodium hydroxide for neutralisation.

(a) Calculate concentration of the:

(i) dilute acid.

(1 mark)

$$C_1 V_1 = C_2 V_2$$

$$0.4 \times 25 = C_2 \times 10$$

$$C_2 = \frac{0.4 \times 25}{10}$$

$$= 1M$$

$$\text{Moles of NaOH} = \frac{0.4 \times 25}{1000} = 0.01$$

$$\text{Moles of HNO}_3 = 0.01$$

$$\text{Molarity of HNO}_3 = \frac{0.01 \times 1000}{10} = 1M$$

(ii) concentrated acid

(1 mark)

$$C_1 V_1 = C_2 V_2$$

$$\frac{1 \times 500}{31.5} = 15.9M$$

$$= 15.9M$$

(b) State the correct method for diluting the concentrated nitric(V) acid.

(1 mark)

Add acid to water ✓

Teacher.co.ke

20. Figure 5 shows part of a radioactive decay series.





Figure 5

(a) Write a nuclear equation for the formation of nuclide K from nuclide X. (1 mark)



PDF Compressor Free Version

- (b) The half-life of nuclide X is 47 minutes. Determine the percentage of nuclide X that remains after 188 minutes. (2 marks)

$$188 = 4 \text{ half-lives} \quad \checkmark 1$$

$$100 \xrightarrow{47} 50 \xrightarrow{47} 25 \xrightarrow{47} 12.5 \xrightarrow{47} 6.25 \quad \checkmark 1$$

$$X \rightarrow \frac{1}{2}X \rightarrow \frac{1}{4}X \rightarrow \frac{1}{8}X \rightarrow \frac{1}{16}X$$

$$\% = \frac{1}{16} \times 100 = 6.25\% \quad \checkmark 1$$

21. Aluminium is extracted from aluminium oxide by electrolysis. (2 marks)
- (a) Other than the cost of electricity, give another reason why this method is expensive. (1 mark)

The graphite anode is replaced periodically.
 has to be replaced.
 $\checkmark 1$

- (b) Calculate the mass of aluminium obtained when a current of 20A is used for 5 hours. (1 Faraday = 96500 C; Al = 27.0) (2 marks)

$Al^{3+} + 3e^- \rightarrow Al$

$Q = It$
 $= 20 \times 5 \times 60 = 360000$

$$\text{moles} = \frac{360000}{3 \times 96500} = 1.244 \text{ moles} \quad \checkmark 1$$

$$\text{mass} = 1.244 \times 27 = 33.588 \text{ g} \quad \checkmark 1$$

22. Explain each of the following observations: (2 marks)

- (a) Articles made of copper turn green when left exposed in air over a long period of time. (1 mark)

Due to formation of copper (II) carbonate.
 Since copper reacts with carbon (IV) oxide.
 $\checkmark 1/2$

(b)

Addition of aqueous ammonia to a solution containing copper(II) ions produces a deep blue solution.

Due to formation of complex ion of tetraammine copper (II) ions $\sqrt{1/2}$

OR due to formation of tetraammine copper (II) ions
 State what is meant by relative atomic mass of an element. $[Cu(NH_3)_4]^{2+}$ (1 mark)
Is the mass of one atom relative to mass of an element compared to the mass of $1/12$ of carbon-12.

23.

(a)


A compound of carbon and element X with formula, CX_n , contains 3.6% carbon by mass. Calculate the relative atomic mass of X. (2 marks)

Let R_{Xm} be n

R _{Xm}	% mass	No. of moles	Ratio
12	3.6	$\frac{3.6}{12}$	1
n	96.4	$\frac{96.4}{n}$	4 $\sqrt{1/2}$
	$\frac{96.4}{n}$	$\frac{96.4}{n}$	$n = 96.4$
			$n = 80.3$

24. Carbon(II) oxide can be prepared by dehydration of ethanedioic acid.

(a) Complete the following equation to show the reaction that takes place. (1 mark)



(b) Name another reagent that can be used to prepare carbon(II) oxide by dehydration. (1 mark)

- Methaneic
- Sodium methanoate

X 002x OF 221 99199 #
18 III 002 II

25. Figure 6 shows an incomplete diagram of a setup for laboratory preparation of nitrogen gas.




Figure 6

- (a) Complete the setup in Figure 6 to show how nitrogen gas can be collected. (2 marks)
- (b) The nitrogen prepared using this setup is purer than that obtained from air. Give a reason. (1 mark)

It has impurities such as noble gases ✓



26.

Hydrazine, $\text{H}-\text{N}-\text{N}-\text{H}$ is used as a fuel in rockets. Using the bond energies in Table 2, calculate the enthalpy change for combustion of hydrazine.



Table 2

Bond	Bond Energy kJ/mol
N—H	388
N—N	163
O=O	496
N≡N	944
O—H	463



(3 marks)

Bonds broken

$$4 \times 388 = 1552$$

$$1 \times 163 = 163$$

$$1 \times 496 = 496$$

$$2211 \checkmark$$

$$\text{Enthalpy of Combustion} = -2796 + 2211$$

$$= -585 \text{ kJ/mol} \checkmark$$

Bonds formed

$$1 \times 944 = 944$$

$$2 \times 463 = 1852$$

$$-2796 \checkmark$$


PDF Compressor Free Version

27. (a) Table 3 gives the standard reduction potentials of some group VII elements.

Table 3

Reduction equations	E°/V
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-$	+1.36
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-$	+1.07
$\text{I}_2 + 2\text{e}^- \rightarrow 2\text{I}^-$	+0.54

State and explain the reactions that take place when aqueous bromine is added to a sample of sea water containing both chloride and iodide ions. (2 marks)



$\text{Br}_2(\text{aq}) + 2\text{Cl}^- \rightarrow \text{No reaction}$ Bromine will oxidize iodide ions to iodine since it has more positive E°

- (b) Bromine will not displace chlorine since E° for Cl^- is more positive ✓
 Give a reason why potassium iodide is added to table salt. (1 mark)

Potassium iodide is a source of iodine. Iodine is needed to regulate functioning of thyroid gland ✓

THIS IS THE LAST PRINTED PAGE.

